


JOURNAL ARTICLE

Morbidity and mortality of people who live close to municipal waste landfills: a multisite cohort study

Francesca Mataloni , Chiara Badaloni, Martina Nicole Golini, Andrea Bolignano, Simone Bucci, Roberto Sozzi, Francesco Forastiere, Marina Davoli, Carla Ancona

International Journal of Epidemiology, Volume 45, Issue 3, June 2016, Pages 806–815, <https://doi.org/10.1093/ije/dyw052>

Published: 24 May 2016 **Article history** ▼

Abstract

Background: The evidence on the health effects related to residing close to landfills is controversial. Nine landfills for municipal waste have been operating in the Lazio region (Central Italy) for several decades. We evaluated the potential health effects associated with contamination from landfills using the estimated concentration of hydrogen sulphide (H_2S) as exposure.

Methods : A cohort of residents within 5 km of landfills was enrolled (subjects resident on 1 January 1996 and those who subsequently moved into the areas until 2008) and followed for mortality and hospitalizations until 31 December 2012. Assessment of exposure to the landfill (H_2S as a tracer) was performed for each subject at enrolment, using a Lagrangian dispersion model. Information on several confounders was available (gender, age, socioeconomic position, outdoor PM_{10} concentration, and distance from busy roads and

industries). Cox regression analysis was performed [Hazard Ratios (HRs), 95% confidence intervals (CIs)].

Results: The cohort included 242 409 individuals. H₂S exposure was associated with mortality from lung cancer and respiratory diseases (e.g. HR for increment of 1 ng/m³ H₂S: 1.10, 95% CI 1.02–1.19; HR 1.09, 95% CI 1.00–1.19, respectively). There were also associations between H₂S and hospitalization for respiratory diseases (HR = 1.02, 95% CI 1.00–1.03), especially acute respiratory infections among children (0–14 years) (HR = 1.06, 95% CI 1.02–1.11).

Conclusions: Exposure to H₂S, a tracer of airborne contamination from landfills, was associated with lung cancer mortality as well as with mortality and morbidity for respiratory diseases. The link with respiratory disease is plausible and coherent with previous studies, whereas the association with lung cancer deserves confirmation.

Keywords: [waste](#), [landfills](#), [residential cohort study](#)

Topic: [child](#), [internship and residency](#), [italy](#), [respiration disorders](#), [respiratory tract infections](#), [socioeconomic factors](#), [hydrogen sulfide](#), [morbidity](#), [mortality](#), [gender](#), [lung cancer](#), [medical residencies](#)

Issue Section: [Environmental Exposures and Cancer](#)

Key Messages

- The evidence on the health of people living close to landfills is still controversial; most of the published studies are characterized by poor exposure assessment, use of health data at the aggregate level and limited possibility of adjusting for socioeconomic status.
- We evaluated the potential health effect of living near nine landfills (Lazio region, Italy), using a residential cohort approach and a dispersion model for exposure assessment.
- Exposure to landfills was associated with mortality from lung cancer and respiratory diseases and with hospitalizations for respiratory diseases, both in adults and in children.

Introduction

People who live close to municipal solid waste (MSW) landfills could be exposed to air pollutants emitted by the plants (landfill gas containing methane, carbon dioxide, hydrogen sulphide and other contaminants including volatile organic compounds, particulate matter and bioaerosols) or to contaminated soil and water. The possible health effects related to residence close to these sites have been assessed in several original papers¹⁻⁹ and evaluated in systematic reviews.^{10,11} Excess of mortality for some cancer sites (e.g. liver, pancreas, kidney, larynx) and non-Hodgkin lymphoma has been noted in some studies,¹⁻³ but the results have not been confirmed in other investigations.⁴⁻⁶ In addition, some studies have indicated an increase of respiratory symptoms among residents close to biodegradable waste facilities.¹² In 2009, Porta

*et al.*¹⁰ concluded that evidence of an association between living close to a landfill and adverse health effects is inconclusive. Most of the published studies have methodological problems, including poor exposure assessment based only on distance from the source, use of health data at the aggregate level and limited possibility of adjusting for socioeconomic status. The quality of the epidemiological studies and scientific knowledge about the issue would be improved by using a residential cohort approach¹³ and applying dispersion models to provide a better exposure assessment.¹⁴

This study aimed at evaluating the association between estimated exposure to hydrogen sulphide (H₂S, produced by anaerobic decomposition of sulphur-containing organic matter in landfills) and mortality and morbidity of a cohort of residents living within 5 km of the nine MSW landfills of the Lazio region (Central Italy, about 5 million inhabitants including the city of Rome). The study was part of a larger project on the characteristics of municipal solid waste treatment plants, their emissions and potential health effects in Lazio (www.eraslazio.it).

Methods

Study areas

Nine municipal solid waste landfills have been operating in Lazio for several decades. Only in the past two decades they were equipped with containments (including leachate collection and treatment, landfill cap construction and landfill gas collection and treatment). The main characteristics of the landfills (together with other potentially relevant environmental factors in the areas, e.g. arsenic contamination)¹⁴ are described in [Supplementary Table 1](#), Landfill characteristics, and in [Supplementary Figure 2](#), Study

areas, (available as [Supplementary data](#) at *IJE* online). The study area was defined for each landfill as a 5-km radius from the boundary of the landfills assessed using GIS software and regional technical maps with a scale of 1:5000. The World Geodetic System of 1984, with the Universal Transverse Mercator zone 33Nord projection (WGS84_UTM33N) was the reference for the geographical coordinates.

Volume 45, Issue 3

June 2016

Exposure assessment

Article Contents

Abstract H₂S has been considered a surrogate measure of all contaminants emitted by landfills, and the airborne concentrations were predicted using a dispersion model. Dispersion models, such as the one we have been using here, have been recently used to assess the health effects of waste management processes.¹⁵⁻¹⁷ We followed a process in three steps. First, yearly H₂S emissions from each sector of the landfills were estimated using a Landfill Gas Emissions Model.¹⁸ Using several variables (the start and end dates of operations for each sector of the landfills, the waste capacity and waste acceptance rate), the annual emission rates for H₂S were calculated by means of a first-order decomposition rate

References

Supplementary data equation: $Q_{H_2S} = \sum_{t=1}^n \sum_{j=0.1}^1 KL_0 \left(\frac{M_t}{10} \right) e^{-kt_{ij}}$ where:

[< Previous](#) [Next >](#)

Q_{H_2S} = annual emission rate (m³/year)

Oxford University Press uses cookies to enhance your experience on our website. By selecting you are agreeing to our use of cookies. You can change your cookie settings at any time. More can be found in our [Cookie Policy](#).

$n = (\text{year of the calculation}) - (\text{initial year of waste acceptance})$

$j = 0.1$ year time increment

K = hydrogen sulphide generation rate (year⁻¹)

L_o = potential hydrogen sulphide generation capacity (m³/Mg)

M_t = mass of waste accepted until t (in Mg)

t_{ij} = age of the waste mass accepted until the i th year (M_t) at the j th section

Mg = Megagram.

We used inventory defaults parameters derived from the US Environmental Protection Agency (EPA) Compilation of Air Pollutant Emission Factors¹⁹ to define hydrogen sulphide generation rate (K) and potential hydrogen sulphide generation capacity (L_o), and M_t and t_{ij} were defined by the Lazio Environmental Protection Agency (EPA) using local data. Second, the EMMA software was used for the temporal and spatial modulation of the estimated emissions. EMMA approximates landfills shape as a regular grid with a resolution of 125 m x 125 m.²⁰ Finally, we used a Lagrangian particle model (SPRAY ver.5, ARIANET Srl, Italy) to simulate H₂S concentrations around the landfills and to produce maps of annual average concentrations around the sites; 2008 was chosen as the reference year for all the sites. The meteorological data were derived from regional measurements made by Lazio EPA in 2005 (that year is considered representative of the meteorological conditions in the area), and used in connection with RAMS data.²¹ The Lagrangian model simulates the transport, dispersion and deposition of pollutants emitted using the orography, the meteorological data, the turbulence and the hourly spatial distribution (horizontal and vertical) of the emissions, based on the characteristics of the single source and on the mass fluxes. The model follows the path of fictitious particles in the atmospheric turbulent flow, and it is able to take into account complex situations, such as the presence of obstacles, breeze cycles, strong meteorological non-

homogeneities and non-stationary, calm wind conditions.

Each subject in the cohort (see below) was assigned an H₂S exposure value corresponding to the estimated annual average value from the dispersion model at the baseline address. In other words, no exposure variation over time was considered and each person remained at the same exposure level during the all study period.

Enrolment of the cohort and follow-up procedures

All residents living within 5 km of the borders of the landfill on 1 January 1996, or those who later moved to the areas until 31 December 2008, were enrolled; datasets from 16 municipalities were used. Vital status was assessed using local registries until 31 December 2012. We considered subjects at risk until they died or moved out of the municipality.

Health outcomes

We analysed natural and cause-specific mortality and hospital admissions for cardiorespiratory diseases. The underlying cause of death for deceased subjects was retrieved from the Regional Registry of Causes of Death, and hospital admissions were obtained from the Regional Hospital Information System which collects information related to all hospital admissions that occur each year in public and private hospitals. Causes of death and diagnoses of hospitalization were coded according to the ICD 9 revision. For each subject, only the principal diagnosis that was the reason for the hospitalization was used and the event (i.e. failure in the Cox model) was defined at the time of the first hospitalization for a specific cause that occurred in the study period. Respiratory hospital admissions for children (residents under 14 years) were

also analysed.

Covariates

We considered for each subject an area-based socioeconomic position (SEP) index, based on several characteristics at the census tract level (around 400 inhabitants) such as education level, occupation, housing conditions, family size and country of origin, classified into five levels (high, middle-high, medium, middle-low, low).²² Modelled outdoor PM₁₀ concentrations ($\mu\text{g}/\text{m}^3$) from primary emissions were assigned to the residential addresses of the cohort participants as a measure of background air quality.²³ The dispersion model was based on the integration between the meteorological Regional Atmospheric Modelling System²¹ and the Eulerian Flexible Air Quality Regional Model (FARM, ARIANET Srl, Italy). As an additional indicator of long-term exposure to traffic-related air pollution at the baseline address, we used the Functional Road Class (FRC) (included in the TeleAtlasMultiNet road network) to classify the type of street: motorway (FRC = 0) and major traffic roads (FRC = 1–5). Presence of an industrial plant in the 2-km buffer from the residence was also considered. Information on individual lifestyle factors was not available.

Statistical analysis

The association between landfill H₂S exposure and mortality and hospital admissions was evaluated using Cox proportional hazard regression models [hazard ratios (HRs), 95% confidence intervals (CIs)], with age as the underlying time variable.

For mortality we defined a latency period of 5 years; therefore we considered all cohort participants who were residents of the area on 1 January 1996 (and started the follow-up on 1 January 2001) and those who subsequently moved to the area up until 31

December 2003 (starting the follow-up 5 years after enrolment). No latency was allowed for the analyses of cardiorespiratory hospitalizations. We first compared the mortality and hospitalization risk of residents according to quartiles of the H_2S distribution. We then considered H_2S as a continuous variable, using the value of the annual mean exposure at residence. A linear association was estimated for increments equal to 1 ng/m^3 of H_2S . We considered as potential confounders socioeconomic position (SEP), PM_{10} background concentrations, residence within 150 m of main roads, 500 m from highways and within 1 or 2 km of industrial plants. With the exception of PM_{10} , which was a continuous variable, all other covariates were considered in the model as categorical variables. In addition, the analyses were performed stratifying in the Cox analysis by landfill sites, to take into account the possible different background rates in the various local areas, by gender and by calendar period (1996–2000, 2001–04, 2005–08, 2009–12), to take into account possible time-related changes in background rates of mortality and hospitalization. Diagnostic tools were used to check the proportional-hazard assumption for all categorical covariates. If any variable in the individual cohort models violated this assumption, effect estimates were compared with a stratified Cox analysis for that covariate. SAS (SAS Institute, NC) and STATA ver. 12 (StataCorp, TX) software programs were used for the statistical analyses.

Results

A total of 242 409 individuals were enrolled in the cohort from 1996 to 2008 (50.4% females), and H_2S concentrations were estimated for each of them at the address of recruitment. The annual average H_2S exposure levels of the population was rather low, 6.3 ng/m^3 [standard deviation (SD) 22.5]; as expected, people living close to the larger landfills (Latina and Rome) had higher H_2S

S exposure levels [mean = 32.7 ng/m³ (SD 76.3) and mean = 45.8 ng/m³ (SD 59), respectively].

The main characteristics of the study cohort according to H₂S concentrations (divided by quartiles of exposure) are described in [Table 1](#). The distribution of gender, age and vital status was rather similar across exposure categories. However, people living in areas with higher concentrations of H₂S were more likely to be of lower SEP compared with people living in areas with lower exposure. PM₁₀ background concentrations were higher in the most exposed group compared with those in the low exposure category. People in the higher exposure category tended to live farther from high traffic roads (500 m) but closer to highways and industrial plants (0–1 km). There was a good correlation between distance from landfill and H₂S exposure.

Table 1. Descriptive individual and environmental characteristics of the cohort members by hydrogen sulphide (H₂S) exposure

	Total		H ₂ S exposure levels (ng/m ³)					
			<25° perc (<0.77)		25°–50° perc (0.77– 2.1)		50°–75° perc (2 4.2)	
	No.	%	No.	%	No.	%	No.	
Total	242	100	60	100.0	60	100	63	
	409		927		775		962	
Gender								
Males	120	49.6	29	49.0	30	49.6	31	
	232		781		137		979	
Females	122	50.4	31	51.0	30	50.4	31	
	177		146		638		983	

Vital status							
Alive	183 060	75.5	48 306	79.3	45 948	75.6	44 673
Migrant	40 740	16.8	8 169	13.4	10 228	16.8	14 446
Dead	18 609	7.7	4 452	7.3	4 599	7.6	4 843
Age at recruitment (years)							
0–14	53 082	21.9	12 246	20.0	13 011	21.4	16 266
15–44	112 754	46.5	27 380	45.0	28 383	46.7	30 661
45–64	50 146	20.7	13 296	22.0	12 584	20.7	11 727
>65	26 427	10.9	8 005	13.0	6 797	11.2	5 308
Area-based socioeconomic position							
High	23 589	9.7	10 012	16.0	6 033	9.9	4 779
Middle-high	41 955	17.3	7 843	13.0	8 834	14.5	9 548
Medium	42 286	17.4	7 447	12.0	8 588	14.1	13 958
Middle-low	50 394	20.8	5 364	9.0	16 816	27.7	17 563
Low	62	25.6	22	37.0	15	25.0	11

	157		806		206		906
Missing	22 028	9.1	7 455	12.0	5 298	8.7	6 208
PM₁₀ (µg/m³)							
< 11.99 (<50° perc)	121 222	50.0	44 371	73.0	29 696	48.9	23 986
11.99–17.69 (50°–90° perc)	96 369	39.8	16 556	27.0	28 967	47.7	31 661
> 17.69 (>90° perc)	24 818	10.2	0	0.0	2 112	3.5	8 315
Distance from major roads (metres)							
<= 150 m	114 698	47.3	31 842	52.0	25 876	42.6	34 506
> 150 m	127 711	52.7	29 085	48.0	34 899	57.4	29 456
Distance from highways (metres)							
<= 500 m	9 428	3.9	2 908	5.0	1 087	1.8	744
> 500 m	232 981	96.1	58 019	95.0	59 688	98.2	63 218
Distance from industrial plants (km)							
0–1 km	12 863	5.3	376	1.0	2 676	4.4	1 130
1–2 km	50 503	20.8	1 138	2.0	9 589	15.8	28 809

> 2 km	179 043	73.9	59 413	98.0	48 510	79.8	34 023
Distance from landfill (km)							
0–1 km	5 187	2.1	0	0.0	3	0.0	19
1–2 km	21 475	8.9	2	0.0	4 225	7.0	5 835
2–3 km	65 386	27.0	8 372	13.7	20 588	33.9	23 627
3–4 km	77 722	32.1	19 739	32.4	18 787	30.9	20 217
4–5 km	72 639	30.0	32 814	53.9	17 172	28.3	14 264

At the end of the follow-up there were 18 609 deaths (7.7%), and for 40 740 subjects (16.8%) the follow-up ended at the time of move away from the municipality of residence.

Table 2 shows the association between H₂S concentrations and cause-specific mortality; effect estimates are given for the quartile distribution of H₂S (25–50, 50–75 and > 75 percentile of the distribution vs < 25 percentile) and for a linear increase of H₂S equal to 1 ng/m³. There were associations between H₂S exposure and lung cancer (HR 1.34, 95% CI 1.06–1.71), and respiratory diseases (HR 1.30, 95% CI 0.99–1.70) when comparing residents in areas with H₂S concentrations greater than 75 percentiles to the reference group. These findings were confirmed when we consider H₂S exposure as linear (HR 1.10, 95% CI 1.02–1.19 for lung cancer and HR 1.09, 95% CI 1.00–1.19 for respiratory diseases). No other associations were noted.

Table 2. Associations between hydrogen sulphide (H₂S in quartiles and continuous) and cause specific mortality: number of deaths (No.) hazard ratios (HR) and 95% confidence intervals (95% CI)

Cause of death (ICD-9-CM)	H ₂ S concentrations						
	<25 ^a percentile	25°–50° percentile			50°–75° percentile		
	No.	No.	Crude HR	HR	95% CI	No.	Crude HR
Natural causes (001–799)	3701	3 946	0.98	1.01	(0.96– 1.06)	4 254	1.00
All cancers (140–239)	1282	1 307	0.97	0.99	(0.91– 1.08)	1 493	1.03
Stomach (151)	75	88	1.03	0.98	(0.70– 1.37)	108	1.27
Colorectal (153–154,159)	154	170	0.99	1.00	(0.79– 1.27)	176	0.96
Liver (155– 156)	102	89	0.86	0.83	(0.61– 1.13)	106	0.89
Pancreas (157)	68	64	0.92	0.93	(0.64– 1.35)	69	0.92
Larynx (161)	17	15	0.81	0.72	(0.33– 1.56)	11	0.38
Lung (162)	276	281	0.98	1.06	(0.89– 1.27)	360	1.09
Bladder (188)	54	48	0.88	0.89	(0.59– 1.36)	56	1.22
Kidney (189)	36	30	0.76	0.85	(0.51– 1.43)	36	0.87

Brain (191)	23	29	1.26	1.25	(0.70– 2.26)	38	1.63
Lymphatic and haematopoietic tissue (200–208)	108	115	1.03	1.16	(0.87– 1.54)	106	0.94
Cardiovascular diseases (390–459)	1457	1681	1.02	1.05	(0.97– 1.13)	1676	0.96
Ischaemic heart diseases (410–414)	512	570	0.99	1.00	(0.88– 1.14)	574	0.86
Respiratory diseases (460–519)	256	244	0.88	0.92	(0.76– 1.11)	279	1.15
Digestive diseases (520–579)	158	163	0.93	0.97	(0.77– 1.24)	218	1.06
Urinary system diseases (580–599)	58	92	1.49	1.54	(1.08– 2.21)	74	1.25

^a Reference category

Table 3 shows the results for cardiorespiratory hospital admissions. No association was detected for cardiovascular diseases. There was an association between the highest quartile of exposure to H₂S and hospitalizations for respiratory diseases (HR 1.05, 95% CI 0.99–1.11) also when considering H₂S exposure as linear (HR 1.02, 95% CI 1.00–1.03). H₂S exposure was linked with respiratory diseases and acute respiratory infection hospital admissions among children (for the highest quartile, HR 1.11, 95% CI 1.01–1.22; HR 1.20, 95% CI 1.04–1.38, respectively) also when we

considered H₂S exposure as a linear term in the model. We found an association with paediatric admissions for asthma but with wider confidence intervals. In both mortality and hospitalization analyses, we did not find effect modification by gender (data not shown).

Table 3. Associations between hydrogen sulphide (H₂S, in quartiles and continuous) and cardiorespiratory morbidity: number of people hospitalized (No.), hazard ratios (HR) and 95% confidence intervals (95% CI)

Diagnosis (ICD-9-CM)	H ₂ S concentrations						
	<25 ^a percentile	25°–50° percentile			50°–75° percentile		
	No.	No.	Crude HR	HR	95% CI	No.	Cru HR
<i>Total cohort</i>							
Cardiovascular diseases (390–459)	6666	6090	0.99	0.99	(0.95–1.03)	6291	0.99
Cardiac diseases (390–429)	3991	3585	0.99	0.98	(0.93–1.03)	3580	0.97
Ischaemic heart diseases (410–414)	1393	1347	1.03	1.02	(0.94–1.10)	1288	0.94
Cerebrovascular diseases (430–438)	1635	1482	0.98	0.98	(0.91–1.06)	1466	0.97
Respiratory diseases (460–	4372	4249	0.98	0.97	(0.92–1.01)	5628	1.02

519)							
Acute respiratory infections (460–466,480–487)	1447	1 441	1.00	0.96	(0.89– 1.04)	1 721	1.00
COPD (490–492;494;496)	654	592	0.96	0.94	(0.84– 1.06)	535	0.92
Asthma (493)	332	355	1.01	1.00	(0.86– 1.17)	594	1.16
<i>Children 0 – 14 years old</i>							
Respiratory diseases (460–519)	1457	1 522	1.00	0.99	(0.92– 1.07)	2 420	1.07
Acute respiratory infections (460–466,480–487)	573	669	1.10	1.02	(0.91– 1.15)	925	1.15
Asthma (493)	257	267	0.98	0.99	(0.83– 1.19)	506	1.23

^a Reference category

Because of the peculiarity of the urban site in Rome ('Malagrotta') (where a large landfill, an incinerator of medical wastes, and a petrochemical refinery are located within just a few kilometres of each other³), we repeated the analyses excluding the subjects who live close to the Malagrotta landfill. There were no important changes in the results (See [Supplementary Tables 3](#) 'Mortality excluding Malagrotta landfill' and [4](#) 'Morbidity excluding Malagrotta landfill', available as [Supplementary data](#) at *IJE* online). We did perform the same sensitivity analysis excluding each landfill at the time, and again the results were similar (see

[Supplementary Figures 7](#) ‘Lung cancer mortality’, [8](#) ‘Respiratory mortality’, [9](#) ‘Respiratory morbidity’ and [10](#) ‘Respiratory morbidity in children’, available as [Supplementary data](#) at *IJE* online).

An additional analysis was performed using distance from the landfills (0–2 km, 2–3 km vs 3–5 km), instead of estimated H₂S concentration, as the exposure variable. Although the results for mortality using distance were not similar to what has been observed using H₂S concentrations (see [Supplementary Table 5](#) ‘Mortality by distance’, available as [Supplementary data](#) at *IJE* online) the results for hospitalizations were similar to those obtained using H₂S concentrations (see [Supplementary Table 6](#) ‘Morbidity by distance’, available as [Supplementary data](#) at *IJE* online).

Our final concern was that migration outside the areas could bias the results in the case of migration being associated with the exposure and if residents with pre-existing diseases were more likely to migrate. We compared the characteristics of people who migrated outside the study areas (40 740 subjects) with those who remained in the areas until the end of the follow-up (201 669 subjects) See [Supplementary Table 11](#) ‘Comparison between migrant and not migrant’, available as [Supplementary data](#) at *IJE* online). We considered gender, age, socioeconomic status and H₂S exposure as fixed variables. Since occurrence of hospitalizations before migration is a time-dependent variable, we compared subjects migrating in the period 2004–12 (19 695 subjects) with all subjects who did not migrate before that period (189 560 subjects), evaluating the occurrence of cardiorespiratory hospitalizations during 1998–2003. Migration was associated with male gender, younger age and lower exposure to H₂S; no clear differences of migrants compared with non-migrants were found for socioeconomic status. In a multinomial logistic regression (data not shown), we found no major differences between the two

groups for respiratory diseases, whereas migrants were less likely than non-migrants to suffer from two or more hospitalizations for cardiovascular disease (OR, 0.74, 95% CI 0.57–0.95) before migrating. All these results indicate that bias due to increased susceptibility of migrants is unlikely given that migrants are less exposed and tend to be healthier than non-migrants.

Discussion

We found a positive association between exposure to hydrogen sulphide (H_2S), that we used as a surrogate for all the pollutants co-emitted from the landfills, and mortality for lung cancer and respiratory diseases as well as hospital admissions for respiratory diseases, especially in children.

Previous studies have investigated the association between residence close to landfills and cancer incidence or cause-specific mortality, with conflicting results. A Canadian cohort study compared cancer incidence in males living close to a landfill with that of residents of farther away areas.¹ The distance from the landfill was assigned to each person based on the residential address at diagnosis. Excess risks for non-Hodgkin lymphoma and liver, pancreas and kidney cancers were found in male residents close to the site. Malagrotta (Rome) residents who lived near (in an area about 2 km²) a large landfill of municipal solid waste, an incinerator and a petrochemical refinery showed an association between proximity to landfill and laryngeal cancer.² A more recent residential cohort study of the same area found that H_2S exposure from the landfill was related to higher risk of mortality from laryngeal cancer and bladder cancer in women, as well as hospitalizations for cardiorespiratory diseases.³ Jarup *et al.* compared cancer incidence (bladder, brain and hepatobiliary cancers and leukaemias) in the population resident within 2 km of

9565 landfills in UK with cancer rates of those who lived more than 2 km away.⁴ Despite the large statistical power, the study did not show excess cancer risk associated with proximity to landfill sites. An ecological study compared mortality, hospital admissions and reproductive health of a population living near a landfill site in Wales with another population matched for socioeconomic status.⁵ No differences between the two populations were found. A study in Brazil evaluated the association between residence close to solid waste landfill sites and cancer mortality.⁶ The exposed areas were defined using a 2-km buffer radius around 15 sites. Standardized mortality ratios were analysed in Bayesian spatial models. The results did not indicate any excess risk for people close to landfills. Some elevated risks of bladder and liver cancer, and death due to congenital malformation were found, although they did not have statistical significance.

The results we found regarding respiratory diseases are consistent with others suggesting a relationship between living close to landfill areas and damage to the respiratory system,^{24,25} as highlighted in a recent systematic review.²⁶ Occurrence of respiratory symptoms was documented among residents living close to waste sites¹² and was linked to inhalation exposure to endotoxin, microorganisms, and aerosols from waste collection and land filling.²⁷

Occupational exposure to organic dust, particulate matters from microbial, plant or animal origin, has been associated with an increased risk of lung cancer in a pooled analysis of case-control studies.²⁵ High lung cancer mortality was found among male residents of Italian National Priority Contaminated Sites with industrial waste landfills or illegal dumps²⁹ and among residents living near incinerators and landfills of hazardous waste in Spain,³⁰ but the overall evidence that residing near landfills is associated with increased risk of lung cancer is still inadequate.¹⁰

This study attempted to overcome some of the limitations of the previously conducted studies, which included issues of study design, exposure assessment and confounding.¹¹ We used a residential cohort approach to provide a more detailed estimation of the population at risk. To each subject in the cohort we assigned an H₂S exposure value (corresponding to the estimated H₂S concentration at the baseline address). It was not possible to consider indexes of average or cumulative exposure based on the different residences, because only a few municipality databases provided information about changes of residence during the follow-up. For this reason, individual exposure reflects residence at the beginning of the follow-up.

Previous studies have considered distance from landfills as a proxy of exposure.^{4,7,9} Distance-to source is easy to understand because it assumes that people living near the landfill are more exposed than people living further away. We used modelled H₂S concentrations as an exposure measure of the landfill gases, on the assumption that the pollution from landfills does not spread uniformly around the site but depends on the quantity of incoming waste, the prevailing winds and the orography of the area.³ Our results for hospitalizations were confirmed when we used distance from the source as the exposure variable instead of modelled H₂S concentrations. There are, however, several aspects in the exposure assessment process we used that should be considered. H₂S generation rates were taken from EPA published material, and waste acceptance capacity and waste acceptance rates were derived from legal authorized values. It is likely, then, that the derived absolute emissions data were more accurate for the recent period and less certain for the past. On the other hand, we used the shape of the H₂S concentrations on the ground to rank subjects as more exposed or less exposed, and this shape is of greater importance than the exact absolute values. Of course, the major limitation of our exposure assessment is related to the lack of a

validation study with *in situ* measurements. Nonetheless, SPRAY is a consolidated model that has been validated using a ‘conventional’ validation framework,³¹ and its performances and efficiency have been evaluated and validated in multiple real conditions with different orography, size of domain, number of grid cells in the domain, meteorological conditions and emission types.^{32–34} The model has been already used in other locations to study health effects of waste management.^{3,17} Another aspect of concern is the use of meteorological parameters that greatly influence the dispersion of the pollutants. We considered the year 2005 as representative of the study area meteorological conditions because there were no particular meteorological anomalies in that year. Running the dispersion model with meteorological data for different years could change the landfills footprint only in presence of extreme weather conditions that strongly affect the annual average. In our opinion, the difference among years is generally minimal and the uncertainty associated with the use of specific meteorological data is negligible.

Our results were adjusted for several confounders: age, socioeconomic position and variables related to the environmental context (proximity to roads with heavy traffic, proximity to industrial sites, air quality) that might otherwise distort the study association. In particular, high level of PM₁₀ (> 90 percentile of the distribution vs < 50 percentile) was associated in our model with cardiovascular and respiratory hospitalizations (HR 1.08, 95% CI 1.01–1.16 and HR 1.03, 95% CI 0.96–1.12, respectively). However, no data were available on the personal habits of the subjects, which could have had a role in the diseases investigated, especially cigarette smoking but also alcohol use, physical activity and obesity. The collection of this information, through telephone interviews or home visits, would have been prohibitive for such a large cohort, and the lack of this information may have biased the results because of confounding not controlled in the analysis. It

should be noted, however, that many personal habits are associated with socioeconomic position. It is therefore reasonable to assume that the analysis that adjusted for socioeconomic index also took into account others individual variables, including smoking. Moreover, excess of hospitalizations for respiratory diseases were found also in children, and no excess mortality/morbidity for cardiovascular diseases (indicative of most of the unmeasured lifestyle factors including smoking) was found, despite the larger statistical power than for respiratory diseases. Therefore, although residual confounding cannot be excluded, it is unlikely that the observed relationship between H₂S exposure and respiratory disturbances could be entirely due to unmeasured smoking habits and other factors.

In conclusion, we found associations between H₂S exposure from landfills and mortality from lung cancer as well as mortality and morbidity for respiratory diseases. The link with respiratory diseases has been observed in other studies and it is potentially related to irritant gases and other organic contaminants. The excess of lung cancer is a relatively new finding.

Funding

This study was supported by the Lazio Waste General Directorate (DGR n929/08) as a part of a larger project on the health effects of waste treatment plants in the Lazio region (ERAS Lazio: Epidemiology, Waste, Environment and Health — www.eraslazio.it). The funder had no scientific role.

Acknowledgements

We wish to thank Margaret Becker for her help in editing the manuscript and Carlo A. Perucci, former director of our department, for initiating the ERAS Lazio project and for his long-standing support of environmental epidemiology.

Conflict of interest : None declared.

References

- 1 Goldberg MS Siemiatyck J DeWar R Dèsy M Riberdy H. Risk of developing cancer relative to living near a municipal solid waste landfill site in Montreal, Quebec, Canada . *Arch Environ Health*1999 ; 54 : 291 – 96 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 2 Michelozzi P Fusco D Forastiere F et al. Small area study of mortality among people living near multiple sources of air pollution . *Occup Environ Med*1998 ; 55 : 611 – 15 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 3 Ancona C Badaloni C Mataloni F et al. . Mortality and morbidity in a population exposed to multiple sources of air pollution: a retrospective cohort study using air dispersion models . *Environ Res*2015 ; 137 : 467 – 74 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 4 Jarup L Briggs D de Hoogh C et al. . Cancer risks in populations living near landfill sites in Great Britain . *Br J Cancer*2002 ; 86 : 1732 – 36 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 5 Fielder HMP Poon-King CM Palmer S R et al. Assessment of impact on health of residents living near the Nant-y-Gwyddon landfill site: retrospective analysis . *BMJ*2000 ; 320 : 19 – 22 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 6 Gouveia N Prado RR. Health risks in areas close to urban solid waste landfill

sites . *Rev Saude Publica*2010 ; 44 : 859 – 66 .

[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)

- 7 Pukkala E Pönkä A. Increased incidence of cancer and asthma in houses built on a former dump area . *Environ Health Perspect*2001 ; 109 : 1121 – 25
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 8 Forastiere F Badaloni C de Hoogh K et al. . Health impact assessment of waste management facilities in three European countries . *Environ Health*2011 ; 10 : 53 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 9 WHO . Population health and waste management: scientific data and available options. <http://www.euro.who.int/document/E91021.pdf> (24 March 2015, date last accessed).
- 10 Porta D Milani S Lazzarino AI et al. Systematic review of epidemiological studies on health effects associated with management of solid waste . *Environ Health*2009 ; 23 : 8 – 60 .
[Google Scholar](#) [WorldCat](#)
- 11 Giusti L. A review of waste management practices and their impact on human health . *Waste Manag*2009 ; 29 : 2227 – 39 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 12 Blanes-Vidal V Bælum J Schwartz J et al. Respiratory and sensory irritation symptoms among residents exposed to low-to-moderate air pollution from biodegradable wastes . *J Expo Sci EnvironEpidemiol*2014 ; 24 : 388 – 97 .
[Google Scholar](#) [Crossref](#) [WorldCat](#)
- 13 Ancona C Mataloni F Badaloni C et al. . Residential cohort approach in industrial contaminated sites: the ERAS Lazio project . *Epidemiol Prev*2014 ; 38(Suppl 1) : 158 – 61 .
[Google Scholar](#) [WorldCat](#)

- 14 D'Ippoliti D Santelli E De Sario M et al. . Arsenic in drinking water and mortality for cancer and chronic diseases in Central Italy, 1990-2010 . *PloS One*2015 ; 10 : e0138182 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 15 Ranzi A Fano V Erspamer L et al. . Mortality and morbidity among people living close to incinerators: a cohort study based on dispersion modeling for exposure assessment . *Environ Health*2011 ; 10 : 22 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 16 Candela S Ranzi A Bonvicini L et al. . Air pollution from incinerators and reproductive outcomes: a multisite study . *Epidemiology*2013 ; 24 : 863 – 70 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 17 Golini MN Ancona C Badaloni CA et al. . Morbidity in a population living close to urban waste incinerator plants in Lazio Region (Central Italy): a retrospective cohort study using a before-after design . *Epidemiol Prev*2014 ; 38 : 323 – 34 .
[Google Scholar](#) [PubMed](#) [WorldCat](#)
- 18 Alexander A Burklin C Singleton A et al. *Landfill Gas Emissions Model (LandGEM) Version 3.02 User's Guide* , May 2005.
- 19 US Environmental Protection Agency . *Compilation of Air Pollutant Emission Factors; AP-42* . <http://www3.epa.gov/ttnchie1/ap42/> (24 March 2015, date last accessed).
- 20 Calori G Radice P. *Emission Manager – Reference Guide* . Milan, Italy : ARIANET R2004.29, 2004 .
[Google Scholar](#) [Google Preview](#) [WorldCat](#) [COPAC](#)
- 21 Pielke RA Cotton WR Walko RL et al. . A comprehensive Meteorological Modeling System RAMS . *Meteorol. Atmos. Phys.*1992 ; 49 : 69 – 91 .
[Google Scholar](#) [Crossref](#) [WorldCat](#)

- 22 Cesaroni G Agabiti N Rosati R Forastiere F Perucci CA. An index of socioeconomic position based on 2001 Census, Rome . *Epidemiol Prev*2006 ; 30 : 352 – 57 .
[Google Scholar](#) [PubMed](#) [WorldCat](#)
- 23 Sozzi R Bolignano A Barberini S Di Giosa AD . apporto sullo stato della qualitell aria nella Regione Lazio 2011. ARPA Lazio. 2012.
- 24 Heaney CD Wing S Campbell RL et al. . Relation between malodor, ambient hydrogen sulfide, and health in a community bordering a landfill . *Environ Res*2011 ; 111 : 847 – 52 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 25 Corrêa CR Abrahão CE CarpinteroMdo C AnarumaFilho F. Landfills as risk factors for respiratory disease in children . *J Pediatr (Rio J)* 2011 ; 87 : 319 – 24 .
[Google Scholar](#) [PubMed](#) [WorldCat](#)
- 26 Mattiello A Chiodini P Bianco E et al. . Health effects associated with the disposal of solid waste in landfills and incinerators in populations living in surrounding areas: a systematic review . *Int J Public Health*2013 ; 58 : 725 – 35 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 27 Park DU Ryu SH Kim SB Yoon CS. An assessment of dust, endotoxin, and microorganism exposure during waste collection and sorting . *J Air Waste Manag Assoc*2011 ; 61 : 461 – 68 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 28 Peters S Kromhout H Olsson AC et al. . Occupational exposure to organic dust increases lung cancer risk in the general population . *Thorax*2012 ; 67 : 111 – 16 .
[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)
- 29 Fazzo L Minichilli F Pirastu R et al. . A meta-analysis of mortality data in Italian contaminated sites with industrial waste landfills or illegal dumps .

*Ann Ist Super Sanità*2014 ; 50 : 278 – 85 .

[Google Scholar](#) [PubMed](#) [WorldCat](#)

- 30 García-Pérez J Fernández-Navarro P Castelló A et al. . Cancer mortality in towns in the vicinity of incinerators and installations for the recovery or disposal of hazardous waste . *Environ Int*2013 ; 51 : 31 – 44 .

[Google Scholar](#) [Crossref](#) [PubMed](#) [WorldCat](#)

- 31 Olesen HR. Toward the establishment of a common framework for model evaluation . In: Gryning SE Schiermeier F (eds). *Air Pollution Modeling and Its Application XI* . New York, NY : Plenum Press , 1996 .

[Google Scholar](#) [Google Preview](#) [WorldCat](#) [COPAC](#)

- 32 Brusasca G Tinarelli G Anfossi D. Particle model simulation of diffusion in low windspeed stable conditions , *Atmos Environ*1989 ; 26 , 707 – 723 .

[Google Scholar](#) [Crossref](#) [WorldCat](#)

- 33 Brusasca G Tinarelli G Anfossi D. Comparison between the results of a Monte Carlo atmospheric diffusion model and tracer experiments , *Atmosp Environ*1992 ; 23 : 1263 – 80 .

[Google Scholar](#) [Crossref](#) [WorldCat](#)

- 34 Anfossi D Tinarelli G TriniCastelli S et al. . A new Lagrangian particle model for the simulation of dense gas dispersion . *Atmos Environ*2010 ; 44:753 – 62 .

[Google Scholar](#) [Crossref](#) [WorldCat](#)

© The Author 2016. Published by Oxford University Press on behalf of the International Epidemiological Association.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work properly cited. For

commercial re-use, please contact journals.permissions@oup.com

Supplementary data

[Supplementary Data](#) - zip file